

Promoting HIV testing in primary care following a randomised trial: An MRC phase IV study

Blizard Institute, Centre for Primary Care and Public Health

Werner Leber¹, Farah El-Shogri¹, Lee Beresford^{1,1}, Estela Capelas Barbosa², Heather McMullen¹, Kambiz Boomla¹, Alison Brown³, Sam Pelluri⁴, Maryam Shahmanesh², Claudia Estcourt^{5,6}, Jane Hutchinson⁶, Danna Millett⁴, Sarah Creighton⁴, Jose Figueroa⁷, Jane Anderson⁴, Steve Morris², Naomi Fulop², Chris Griffiths¹

¹Queen Mary, University of London; ²University College London; ³Public Health England; ⁴Homerton University Hospital NHS Foundation Trust; ⁵Glasgow Caledonian University; ⁶Barts Health NHS Trust; ⁷NHS England & NHS City and Hackney

Background

HIV remains underdiagnosed. Guidelines recommend routine HIV testing in primary care but evidence on implementing testing is lacking.¹⁻³

In the RHIVA2 randomised controlled trial, we showed that integrating HIV screening using rapid point-of-care HIV testing into general practice registration health checks in Hackney, a deprived inner London borough with high HIV prevalence, delivered increased and early diagnosis of HIV.⁴

However, interventions effective in the context of a trial may be less so when implemented in routine practice.

Following the RHIVA2 trial, we wished to implement HIV testing across the borough. 20 practices had received HIV testing training during the trial. Thus of the 44 practices in Hackney, we attempted to: a) implement training in the 24 naïve practices (20 control, 4 non-participating), and b) reinforce training in the 20 intervention practices.

Objectives

To determine the effectiveness of the RHIVA2 intervention when implemented as a routine clinical service outside a trial setting.

Study design

MRC phase IV study, using an interrupted time series analysis.

Study duration

19 April, 2010 to 30 June, 2015.

Study population

- Individuals aged 16 years or older registering at study practices
- Individuals able to undertake the pre-test discussion in English or with a suitable translator.

RHIVA2 Interventions

The RHIVA2 trial programme (2010-2012) included:

- Practice-based educational training session for the primary care team to promote rapid HIV testing (INSTI™ HIV-1/HIV-2 Rapid Antibody Test, bioLytical Laboratories, Canada) at GP registration
- Follow up meeting with a nominated practice lead nurse
- Incentive payment of £10 per rapid HIV test performed
- External quality assessment
- Regular data monitoring for safe diagnosis and referral of newly diagnosed patients into secondary care.

The RHIVA2 post-trial implementation package (2012-2015) consisted of:

- Additional modified training to offer both rapid and serology HIV testing at all practices in any clinical setting
- Service evaluation of antiretroviral prescribing and missed opportunities of HIV diagnosis in primary care⁶
- Addition of incentive payments between £7 and £10 per HIV test performed to the sexual health local enhanced service.

Data collection

Electronic patient record searches for surgery recruitment data, patient demographics and HIV test results.

Study findings

For HIV promotional activities and testing analysis, please see Figures 1 & 2.

Implementation uptake:

A total of 12 naïve practices (11 former trial control practices and 1 former non-participating practice) received the implementation package, and 6 former trial intervention practices were reinforced; 31 practices (13 former intervention, 15 former control, and 3 former non-participating practices) completed the service evaluation.

Immediate increase in HIV testing:

Amongst practices that received the RHIVA2 intervention, testing rates increased immediately following the intervention by 85% (IRR = 1.85, 95% CI 1.72 to 1.99), and a similar effect (IRR 1.82, 95% CI 1.72 to 1.93) was observed when including the comparator practices that had not received the intervention during the trial period.

Gradual decline in HIV testing over time:

The change in the effect of the intervention following the training phase decreased minimally, albeit significantly, over time (IRR = 0.994, 95% CI 0.990 to 0.997).

Conclusions

Implementation of HIV testing through research significantly increased HIV testing rates, but regular re-training may be required to keep it sustainable long term.

Figure 1: Sexual health service provision and research in Hackney





Figure 2: Increased HIV testing in Hackney general practice



The medium blue line shows the effect on testing of training 20 trial intervention practices; the dark blue line shows the effect on HIV testing of training 12 naïve practices; the light blue line shows testing in naïve practices that declined training. Graphs are shown as if all practices were trained at the same time.

Funding

NIHR Collaborations for Leadership in Applied Health Research and Care (CLAHRC), NHS City and Hackney, Department of Health.

Acknowledgements

We thank all participants and general practices. In addition, we would like to thank the following people: Keith Prescott, Martin Sharp, and Jack Dunne.

References

1. British HIV Association. UK National Guidelines for HIV testing. 2008.
2. ECDC. HIV testing: increasing uptake and effectiveness in the European Union. 2014.
3. National Institute for Health and Care Excellence. HIV testing: increasing uptake among people who may have undiagnosed HIV (Joint NICE and Public Health England guideline). 2016.
4. Leber W, McMullen H, Anderson J, et al. Promotion of rapid testing for HIV in primary care (RHIVA2): a cluster-randomised controlled trial. *The Lancet HIV*. 2015.
5. Prost A, Griffiths CJ, Anderson J, et al. Feasibility and acceptability of offering rapid HIV tests to patients registering with primary care in London (UK): a pilot study. *Sex Transm Infect*. 2009.
6. Wellesley R, Whittle A, Figueroa J, et al. Does general practice deliver safe primary care to people living with HIV? A case-notes review. *Br J Gen Pract*. 2015.

Contact: w.leber@qmul.ac.uk